

AD-A103 870

WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER

F/G 12/1

ON THE HAAR AND WALSH SYSTEMS ON A TRIANGLE.(U)


JUL 81 Y Y FENG, D. X GI

DAA629-80-C-0041

UNCLASSIFIED

MRC-TSR-2235

NL



AD A103870

MRC Technical Summary Report #2235 ✓

ON THE HAAR AND WALSH SYSTEMS
ON A TRIANGLE

Y. Y. Feng and D. X. Qi

Mathematics Research Center
University of Wisconsin-Madison
610 Walnut Street
Madison, Wisconsin 53706

July 1981

(Received May 26, 1981)

DTIC FILE COPY

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

Approved for public release
Distribution unlimited

81 9 08 036

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

ON THE HAAR AND WALSH SYSTEMS ON A TRIANGLE

Y. Y. Feng^{*} and D. X. Qi^{**}

Technical Summary Report #2235
July 1981

ABSTRACT

In this paper we establish the Haar and Walsh systems on a triangle. These systems are complete in $L_2(\Delta)$. The uniform convergence of the Haar-Fourier series and the uniform convergence by group of the Walsh-Fourier series for any continuous function are proved.

AMS (MOS) Subject Classification: 41A15

Key Words: Haar function, Walsh function, Orthogonal system, Series expansion, Uniform convergence, Triangular domain

Work Unit Number 3 (Numerical Analysis and Computer Science)

^{*}Department of Mathematics, China University of Science and Technology, Hefei, China.

^{**}Department of Mathematics, Jilin University, Changchun, China.

SIGNIFICANCE AND EXPLANATION

A number of papers have been concerned with developing the theories of discontinuous orthonormal systems and their applications. In particular, the Haar and Walsh systems are presently the most important examples of nonsinusoidal functions, and have proved most useful in communication.

Some authors have studied the properties of approximation from the mathematical point of view. It seems interesting and helpful for both theory and practice to investigate the Haar and Walsh functions for a multivariate setting. In fact, many signals in communications and other functions are of several variables (for instance, TV signals have two space variables and the time variable). If the domain of definition of the system is tensor product, then the existing systems are readily extended to several variables.

The problem is how to construct an orthonormal system on a triangular domain in the plane, or more generally, on a simplex in n -dimensional space. This paper defines the Haar and Walsh system on a triangle domain, proves the orthogonality and completeness in L_2 . Also the uniform convergence for the Haar-Fourier series, uniform convergence by group for the Walsh-Fourier series are studied. All of these results can be generalized easily to n dimensions.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Avail and/or	
Dist	Special
A	

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the authors of this report.

ON THE HAAR AND WALSH SYSTEMS ON A TRIANGLE

Y. Y. Feng* and D. X. Qi**

1. Introduction

No doubt, it is interesting and useful to study multivariate Haar and Walsh functions either in theory or in practice. If we investigate on a domain which can be considered a Cartesian product, then the functions are readily extended to several variables from the one variable. Setting by the tensor product construct Harmuth has shown those kinds of multivariate systems in his book [5] and pointed out the applications in communication.

In this paper we attempt to focus on a triangle, or more generally on a simplex in n -dimensional space. We were unable to find any paper about it. Perhaps it puzzles some people temporarily.

The main contribution of this paper is to establish the Haar and Walsh system with two variables on a triangle. We prove their orthonormality and completeness in Hilbert space L_2 . Moreover, the corresponding Haar-Fourier series and Walsh-Fourier series for any continuous function are uniformly convergent and convergent by group respectively.

It is easy to generalize these results to the n -dimensional simplex. For simplicity we will discuss only the two-dimensional triangle.

Now we should explain some preliminaries and notations.

The Haar functions on $[0,1]$ are defined as follows:

$$X_0(t) := 1 \text{ for } 0 \leq t \leq 1,$$

and

*Department of Mathematics, China University of Science and Technology, Hefei, China.

**Department of Mathematics, Jilin University, Changchun, China.

$$\chi_n^{(k)}(t) := \begin{cases} \sqrt{2^n}, & \text{for } \frac{2k-2}{2^{n+1}} < t < \frac{2k-1}{2^{n+1}}, \\ -\sqrt{2^n}, & \text{for } \frac{2k-1}{2^{n+1}} < t < \frac{2k}{2^{n+1}}, \\ 0, & \text{elsewhere in } [0,1] \end{cases} \quad (1.1)$$

$$k = 1, 2, 3, \dots, 2^n; \quad n = 1, 2, 3, \dots, \infty.$$

The Walsh functions on $[0,1]$ consist of the following ones:

$$\begin{aligned} w_0(t) &:= 1 \text{ for } 0 \leq t \leq 1, \\ w_1(t) &:= \begin{cases} 1 & \text{for } 0 \leq t < \frac{1}{2}, \\ -1 & \text{for } \frac{1}{2} < t \leq 1, \end{cases} \\ w_{n+1}^{(2k-1)}(t) &:= \begin{cases} w_n^{(k)}(2t), & \text{for } 0 \leq t < \frac{1}{2}, \\ (-1)^{k+1} w_n^{(k)}(2t-1), & \text{for } \frac{1}{2} < t \leq 1, \end{cases} \\ w_{n+1}^{(2k)}(t) &:= \begin{cases} w_n^{(k)}(2t), & \text{for } 0 \leq t < \frac{1}{2}, \\ (-1)^k w_n^{(k)}(2t-1), & \text{for } \frac{1}{2} < t \leq 1, \end{cases} \\ k &= 1, 2, 3, \dots, 2^n, \quad n = 1, 2, 3, \dots, \infty. \end{aligned} \quad (1.2)$$

Some detailed investigation of the Haar and Walsh systems can be found in [1], [3], [5].

In order to generalize the Haar and Walsh systems to the two-dimensional case we should explain our representation in this paper. The Cartesian coordinates are not very convenient for triangular elements, and a special type of coordinate system called area coordinates should be used.

Referring to Figure 1 it is seen that the internal point P will divide the triangle ABC into three smaller triangles, and depending on the position of the point P , the area of each one of the triangles PAB , PBC , PCA can vary from zero to $|\Delta|$, which is the area of the triangle ABC . In other words, the ratios $\frac{a}{|\Delta|}$, $\frac{b}{|\Delta|}$ and $\frac{c}{|\Delta|}$ will take up any value between zero and unity. Here a , b , c are the area of triangles PBC , PCA , PAB respectively.




Figure 1

These ratios are called area coordinates, and they are defined by $l_1 := \frac{a}{|\Delta|}$,
 $l_2 := \frac{b}{|\Delta|}$, $l_3 := \frac{c}{|\Delta|}$.

It is easy to see that

$$\begin{pmatrix} 1 \\ x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{pmatrix} \begin{pmatrix} l_1 \\ l_2 \\ l_3 \end{pmatrix}.$$

If two points P and Q are in two similar triangles respectively, and have the same area coordinates, then we denote them by $P \sim Q$.

2. An orthonormal sequence χ on a triangular domain

Suppose Δ (or ΔABC) is any triangle on a plane and $|\Delta| = 1$ is the area of the ΔABC . If D, E, F are midpoints of AB, BC, CA respectively, connecting DE, EF, FD , we divide the Δ into four similar small triangles $\Delta ADF, \Delta DBE, \Delta FEC, \Delta EFD$. We call them $\Delta_1, \Delta_2, \Delta_3, \Delta_4$ respectively.

We define the sequence χ as follows:

$$\begin{aligned} \chi_0(P) &:= 1 \text{ for } P \in \Delta, \\ \chi_1^{(1)}(P) &:= \begin{cases} 1 & \text{for } P \in \Delta_1 \cup \Delta_2, \\ -1 & \text{for } P \in \Delta_3 \cup \Delta_4, \end{cases} \\ \chi_1^{(2)}(P) &:= \begin{cases} \sqrt{2} & \text{for } P \in \Delta_1, \\ -\sqrt{2} & \text{for } P \in \Delta_2, \\ 0 & \text{for } P \in \Delta_3 \cup \Delta_4, \end{cases} \\ \chi_1^{(3)}(P) &:= \begin{cases} \sqrt{2} & \text{for } P \in \Delta_3, \\ -\sqrt{2} & \text{for } P \in \Delta_4, \\ 0 & \text{for } P \in \Delta_1 \cup \Delta_2, \end{cases} \\ &\dots \dots \dots \\ \chi_n^{(3j+1)}(P) &:= \begin{cases} 2\chi_{n-1}^{(i)}(Q) & \text{for } P \in \Delta_{j+1}, \\ 0 & \text{for } P \in \Delta \setminus \Delta_{j+1}, \end{cases} \end{aligned} \quad (2.1)$$

where $Q \in \Delta, Q \sim P, j = 0, 1, 2, 3, i = 1, 2, \dots, 3 \cdot 4^{n-2}, n = 2, 3, \dots$.

At a point of discontinuity, let the value of these functions be the average.

Now we consider the orthogonality of the sequence χ . We prove the following theorem.

Theorem 1. The sequence χ defined by (2.1) is orthonormal.

Proof. At first, it is easy to check that when $n \leq 2$ the sequence $\{\chi_n^{(j)}\}$ is orthonormal. We suppose that the theorem holds for $n \leq N$. For $2 \leq m \leq N+1$, $j_1, j_2 = 0, 1, 2, 3; i_1 = 1, 2, \dots, 3 \cdot 4^{N-1}; i_2 = 1, 2, \dots, 3 \cdot 4^{m-2}$, by (2.1) and induction

hypothesis, we get

$$\begin{aligned}
 \int_{\Delta} \chi_{N+1}^{(3j_1+1,1)}(P) \chi_m^{(3j_2+1,2)}(P) dP &= 4\delta_{j_1,j_2} \int_{\Delta_{j_1+1}} \chi_N^{(1,1)}(Q) \chi_{m-1}^{(1,2)}(Q) dQ \\
 &= \delta_{j_1,j_2} \int_{\Delta} \chi_N^{(1,1)}(Q) \chi_{m-1}^{(1,2)}(Q) dQ \\
 &= \delta_{j_1,j_2} \delta_{1,1} \delta_{2,m-1} .
 \end{aligned}$$

It is easy to verify that

$$\int_{\Delta} \chi_{N+1}^{(3j+1,1)}(P) \chi_1^{(1,2)}(P) dP = \int_{\Delta} \chi_{N+1}^{(3j+1,1)}(P) \chi_0(P) dP = 0 .$$

Therefore the theorem holds for $n = N + 1$, and this finishes the induction. ■

3. Convergence properties

The triangle Δ has been divided into four similar smaller triangles Δ_i ($i = 1, 2, 3, 4$). Now set

$$\Delta_{1,i} := \Delta_i \quad (i = 1, 2, 3, 4).$$

For each $\Delta_{1,i}$, we divide it into four similar smaller triangles in the same way as we did before. We order them as $\Delta_{2,1}, \Delta_{2,2}, \dots, \Delta_{2,16}$ such that

$$\Delta_{1,i} = \Delta_{2,4i-3} \cup \Delta_{2,4i-2} \cup \Delta_{2,4i-1} \cup \Delta_{2,4i}, \quad i = 1, 2, 3, 4.$$

We continue this process. For any n we get a sequence $\Delta_{n,1}, \Delta_{n,2}, \dots, \Delta_{n,4^n}$ such that

$$\Delta_{n-1,i} = \Delta_{n,4i-3} \cup \Delta_{n,4i-2} \cup \Delta_{n,4i-1} \cup \Delta_{n,4i}, \quad i = 1, 2, 3, \dots, 4^{n-1}, \quad n = 1, 2, 3, \dots, \infty.$$

$$\Delta_{0,1} := \Delta.$$

Define a function sequence $\{f_{n,i}\}$ on the Δ :

$$\begin{aligned} f_0(P) &:= 1 \quad \text{for } P \in \Delta, \\ f_{1,i}(P) &:= \begin{cases} 1 & \text{for } P \in \Delta_{1,i}, \\ 0 & \text{for } P \in \Delta \setminus \Delta_{1,i}, \end{cases} \quad i = 1, 2, 3, 4, \\ &\dots \dots \dots \dots \\ f_{n,i}(P) &:= \begin{cases} 1 & \text{for } P \in \Delta_{n,i}, \\ 0 & \text{for } P \in \Delta \setminus \Delta_{n,i}, \end{cases} \quad (3.1) \\ &i = 1, 2, 3, \dots, 4^n, \quad n = 1, 2, 3, \dots, \infty. \end{aligned}$$

It is obvious that the sequence $\{f_{n,i}\}$ is orthogonal.

Let

$$M_n := \text{span}(f_{n,1}, f_{n,2}, \dots, f_{n,4^n}) \quad (n \geq 0). \quad (3.2)$$

Thus

$$\dim M_n = 4^n.$$

For convenience, sometimes we use notation

$$\begin{aligned} x_1 &:= x_0, \\ x_{4^{n-1}+i} &:= x_n^{(i)}, \quad n \geq 1, \quad i = 1, 2, \dots, 4^{n-1}. \end{aligned} \quad (3.3)$$

Set

$$H_n := \text{span}(X_1, X_2, \dots, X_n) .$$

It is clear that

$$H_{4^n} = M_n , \quad (3.4)$$

since $H_{4^n} \subset M_n$ and $\dim H_{4^n} = \dim M_n = 4^n$.

We define

$$L_2(\Delta) := \{f \mid \int_{\Delta} f^2 d\sigma < \infty\}$$

and

$$\|f\|_2^2 := \int_{\Delta} f^2 d\sigma .$$

Then the Fourier series of a given function $F \in L_2(\Delta)$ in terms of the function sequence

$\{X_n\}$ is

$$F \sim \sum_{i=1}^{\infty} a_i X_i \quad (3.5)$$

with

$$a_i := \int_{\Delta} F(p) X_i(p) dP .$$

Let

$$P_n F := \sum_{i=1}^n a_i X_i(p) \quad (3.6)$$

be the n -th partial sum of the series (3.5).

From the orthogonality of sequence $\{X_n\}$ we know that $P_n F$ is the best L_2 -approximation to F from H_n . Hence it is convergent to F if F is in $L_2(\Delta)$, since H_n is dense in $L_2(\Delta)$. Thus we get the following theorem.

Theorem 2. If $F \in L_2(\Delta)$, then

$$\lim_{n \rightarrow \infty} \|F - P_n F\|_2 = 0 .$$

In order to study the uniform convergence we let

$$C(\Delta) := \{f | f \text{ is continuous on } \Delta\}$$

and

$$\|f\|_{\infty} := \max_{P \in \Delta} |f(P)|.$$

For $F \in C(\Delta)$ we define

$$P_n^{(j)} F := \int_{\Delta} F X_0 d\sigma \cdot X_0 + \int_{\Delta} F X_1^{(1)} d\sigma \cdot X_1^{(1)} + \dots + \int_{\Delta} F X_n^{(j)} d\sigma \cdot X_n^{(j)}. \quad (3.7)$$

Set

$$\begin{aligned} K_0(P, Q) &:= X_0(P) X_0(Q), \quad (\text{for } P, Q \in \Delta) \\ &\dots \dots \dots \dots \\ K_n^{(j)}(P, Q) &:= X_0(P) X_0(Q) + X_1^{(1)}(P) X_1^{(1)}(Q) + \dots + X_n^{(j)}(P) X_n^{(j)}(Q), \\ &j = 1, 2, \dots, 3 \cdot 4^{n-1}, \quad n = 1, 2, \dots \end{aligned} \quad (3.8)$$

Thus

$$P_n^{(j)} F(P) = \int_{\Delta} K_n^{(j)}(P, Q) F(Q) dQ. \quad (3.9)$$

Let $A := (a_{ij})$ ($i, j = 1, 2, 3, \dots, 4^n$) be any $4^n \times 4^n$ ($n = 1, 2, \dots$) matrix and $G(P, Q)$ be any function defined on $\Delta \times \Delta$.

The notation $G(P, Q) \leftrightarrow A$ means that the value of $G(P, Q)$ is a_{ij} when $P \in \Delta_{n,j}$, $Q \in \Delta_{n,i}$. It leads to the following relationship:

$$\begin{aligned} X_0(P) X_0(Q) \leftrightarrow \sigma_0 &:= \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}, \\ X_1^{(1)}(P) X_1^{(1)}(Q) \leftrightarrow \sigma_1 &:= \begin{pmatrix} 1 & 1 & -1 & -1 \\ 1 & 1 & -1 & -1 \\ -1 & -1 & 1 & 1 \\ -1 & -1 & 1 & 1 \end{pmatrix}, \end{aligned} \quad (3.10)$$

$$\chi_1^{(2)}(P)\chi_1^{(2)}(Q) \leftrightarrow \sigma_2 := \begin{pmatrix} 2 & -2 & 0 & 0 \\ -2 & 2 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

$$\chi_1^{(3)}(P)\chi_1^{(3)}(Q) \leftrightarrow \sigma_3 := \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 2 & -2 \\ 0 & 0 & -2 & 2 \end{pmatrix}.$$

In order to write those more shortly, we use the notation

$$\text{diag block}(A_1, A_2, \dots, A_m) := \begin{pmatrix} A_1 & & & \\ & A_2 & & \\ & & \ddots & \\ 0 & & & A_m \end{pmatrix}, \quad (3.11)$$

where A_i is square submatrix.

Using (3.11) we get

$$\chi_1^{(1)}(P, Q) \leftrightarrow \sigma_0 + \sigma_1 = \text{diag block} \left[\begin{pmatrix} 2 & 2 \\ 2 & 2 \end{pmatrix}, \begin{pmatrix} 2 & 2 \\ 2 & 2 \end{pmatrix} \right],$$

$$\chi_1^{(2)}(P, Q) \leftrightarrow \sigma_0 + \sigma_1 + \sigma_2 = \text{diag block} \left[4, 4, \begin{pmatrix} 2 & 2 \\ 2 & 2 \end{pmatrix} \right],$$

$$\chi_1^{(3)}(P, Q) \leftrightarrow \sigma_0 + \sigma_1 + \sigma_2 + \sigma_3 = \text{diag block}(4, 4, 4, 4),$$

where $\text{diag block}(4, 4, 4, 4) = 4I_4$, I_n is $n \times n$ identity matrix. We denote the $m \times m$ zero-element matrix by O_m below.

Since

$$\chi_2^{(i)}(P)\chi_2^{(i)}(Q) \leftrightarrow \text{diag}(4\sigma_i, O_4, O_4, O_4), \quad (i = 1, 2, 3)$$

we get

$$\kappa_2^{(1)}(P,Q) \leftrightarrow \text{diag block} \left(4 \sum_{j=0}^{\frac{1}{2}} \sigma_j, 4I_4, 4I_4, 4I_4 \right),$$

$$\kappa_2^{(3+1)}(P,Q) \leftrightarrow \text{diag block} \left(4^2 I_4, 4 \sum_{j=0}^{\frac{1}{2}} \sigma_j, 4I_4, 4I_4 \right),$$

$$\kappa_2^{(6+1)}(P,Q) \leftrightarrow \text{diag block} \left(4^2 I_4, 4^2 I_4, 4 \sum_{j=0}^{\frac{1}{2}} \sigma_j, 4I_4 \right),$$

and

$$\kappa_2^{(9+1)}(P,Q) \leftrightarrow \text{diag block} \left(4^2 I_4, 4^2 I_4, 4^2 I_4, 4 \sum_{j=0}^{\frac{1}{2}} \sigma_j \right), \quad (i = 1, 2, 3)$$

especially

$$\kappa_2^{(12)}(P,Q) \leftrightarrow 4^2 I_{16} = \text{diag block} (4^2 I_4, 4^2 I_4, 4^2 I_4, 4^2 I_4).$$

Suppose in the general case that

$$\kappa_n^{(3 \cdot 4^{n-1})}(P,Q) \leftrightarrow 4^n I_{4^n}. \quad (3.12)$$

By definition of (2.1) and (3.8),

$$\kappa_{n+1}^{(L)}(P,Q) \leftrightarrow \text{diag block} (\kappa_{1,1}, \kappa_{2,2}, \dots, \kappa_{4^n, 4^n})$$

where each $\kappa_{1,1}$ is a 4×4 matrix. More precisely

$$\begin{aligned} \kappa_{n+1}^{(1)}(P,Q) &\leftrightarrow \text{diag block} \left(4^n \sum_{t=0}^{\frac{1}{2}} \sigma_t, 4^n I_4, \dots, 4^n I_4 \right), \quad (i = 1, 2, 3), \\ \dots \dots \dots & \\ \kappa_{n+1}^{(3+1)}(P,Q) &\leftrightarrow \text{diag block} \left(4^{n+1} I_4, \dots, 4^n \sum_{t=0}^{\frac{1}{2}} \sigma_t, \dots, 4^n I_4 \right) \end{aligned} \quad (3.13)$$

where the term $4^n \sum_{t=0}^{\frac{1}{2}} \sigma_t$ is the $(j+1)$ -th block, $(j = 1, 2, \dots, 4^n - 1)$ in particular

$$\kappa_{n+1}^{(3 \cdot 4^n)}(P, Q) \rightarrow 4^{n+1} I_{4^{n+1}},$$

therefore for any $n = 1, 2, \dots$ (3.12) holds.

Suppose

$$a \in \Delta_{n, 4(l-1)+i} \subset \Delta_{n-1, l}, \quad i = 1, 2, 3, 4.$$

By (3.9) we know

$$p_n^{(j)} F(a) = \int_{\Delta} \kappa_n^{(j)}(a, Q) F(Q) dQ. \quad (3.14)$$

By (3.13), (3.14) we obtain

$$p_n^{(j)} F(a) = 4^{n-1} \int_{\Delta_{n-1, l}} F(Q) dQ = \frac{1}{|\Delta_{n-1, l}|} \int_{\Delta_{n-1, l}} F(Q) dQ \quad (3.15)$$

for $j \leq 3(l-1)$ and

$$p_n^{(j)} F(a) = 4^n \int_{\Delta_{n, l}} F(Q) dQ = \frac{1}{|\Delta_{n, l}|} \int_{\Delta_{n, l}} F(Q) dQ \quad (3.16)$$

for $j > 3l$.

For $j = 3l-2, 3l-1$ we have

$$p_n^{(3l-2)} F(a) = \frac{2}{|\Delta_{n-1, l}|} \left(\int_{\Delta_{n, 4(l-1)+i}} F(Q) dQ + \int_{\Delta_{n, 4(l-1)+i+1}} F(Q) dQ \right) \quad (3.17)$$

($i = 1$ or 3) and

$$p_n^{(3l-1)} F(a) = \begin{cases} \frac{1}{|\Delta_{n, l}|} \int_{\Delta_{n, 4(l-1)}} F(Q) dQ \\ a \in \Delta_{n, 4(l-1)+i}, \quad i = 1, 2, \\ \frac{2}{|\Delta_{n-1, l}|} \left(\int_{\Delta_{n, 4(l-1)+3}} F(Q) dQ + \int_{\Delta_{n, 4(l-1)+4}} F(Q) dQ \right), \quad i = 3, 4. \end{cases} \quad (3.18)$$

In any case, from (3.15) to (3.18) we conclude

$$\lim_{n \rightarrow \infty} p_n^{(j)} F(a) = \frac{1}{|\Delta_a|} \int_{\Delta_a} F(Q) dQ = F(a), \quad (3.19)$$

where $\Delta_a \in \{\Delta_{n, i}\}$ and $a \in \Delta_a$, $|\Delta_a| \rightarrow 0$ when $n \rightarrow \infty$.

It is easy to check that

$$\int_{\Delta} |\kappa_n^{(j)}(p, Q)| dQ = 1. \quad (3.20)$$

Now (3.19), (3.20) imply the following theorem.

Theorem 3. For $F \in C(\Delta)$ $\lim_{n \rightarrow \infty} |P_n^{(j)} F - F|_{\infty} = 0$ ($j = 1, 2, \dots, 3 \cdot 4^{n-1}$).

4. On the Walsh system

Naturally there exist some different forms of definition which are equivalent. We use area coordinates to define the two-variable Walsh function. Some notations follow the case of the Haar functions.

$$\begin{aligned}
 W_0(P) &:= 1 && \text{for } P \in \Delta; \\
 W_{n+1}^{(i)}(P) &:= W_n^{(i)}(Q) && \text{for } P \in \Delta, \quad i = 1, 2, \dots, 4^n, \\
 W_{n+1}^{(4^n+1)}(P) &:= \begin{cases} \lambda & \text{for } P \in \Delta_1 \cup \Delta_3, \\ -\lambda & \text{for } P \in \Delta_2 \cup \Delta_4, \end{cases} \\
 W_{n+1}^{(2 \cdot 4^n+1)}(P) &:= \begin{cases} \lambda & \text{for } P \in \Delta_1 \cup \Delta_2, \\ -\lambda & \text{for } P \in \Delta_3 \cup \Delta_4, \end{cases} \\
 W_{n+1}^{(3 \cdot 4^n+1)}(P) &:= \begin{cases} \lambda & \text{for } P \in \Delta_1 \cup \Delta_4, \\ -\lambda & \text{for } P \in \Delta_2 \cup \Delta_3, \end{cases} \quad (4.1)
 \end{aligned}$$

where

$$\begin{aligned}
 \lambda &:= W_n^{(i)}(Q), \quad Q \in \Delta, \quad Q \sim P, \quad i = 1, 2, 3, \dots, 4^n, \quad n = 0, 1, 2, \dots \\
 W_0^{(1)}(P) &:= W_0(P) = 1 \quad \text{for } P \in \Delta.
 \end{aligned}$$

This finishes the definition of the sequence W .

Sometimes we prefer W_l ($l = 1, 2, 3, \dots, 4^{m+1}$) to $W_{m+1}^{(j \cdot 4^m + i)}$ with $l = j \cdot 4^m + i$ ($i = 1, 2, \dots, 4^m$, $j = 0, 1, 2, 3$).

At a point of discontinuity, the values of these functions are taken as the average.

Figures 2 and 3 show the Walsh sequence when $n = 0, 1$.

Before the discussion of the orthogonality of the Walsh system we introduce the Hadamard matrix ([2], [4], p. 207).

The Hadamard matrix is a square array whose elements consist only of $+1$ and -1 and whose rows (and columns) are orthogonal to one another. Obviously the lowest order nontrivial Hadamard matrix is of the order two, viz.

$$H_2 := \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}. \quad (4.2)$$

Higher order matrices whose orders are powers of two can be obtained from the recurrent relationship

$$H_n = H_{n/2} \otimes H_2 \quad (4.3)$$

where \otimes denotes the direct or Kronecker product and n is a power of two. The direct product means replacing each element in the matrix by the H_2 . With the help of the Hadamard matrix the one-dimensional Walsh function can be defined [1]. In the two-dimensional case we should use the 4×4 matrix

$$H_4 = H_2 \otimes H_2$$



and get the recurrent relationship

$$H_N = H_{N/4} \otimes H_4. \quad (4.4)$$

The Hadamard matrix (4.4) corresponds to the Walsh sequence $\{w_i\}$ ($i = 1, 2, \dots, 4^N$) for a given N .

Figures 2 and 3 show the Walsh sequence associated with the Hadamard matrix.


In Figures 2 and 3 black areas represent $+1$, white areas -1 . The following triangle shows a certain order.



$$\begin{array}{l}
 n = 0 \\
 \left. \begin{array}{l} w_1^{(1)} \\ w_1^{(2)} \\ w_1^{(3)} \\ w_1^{(4)} \end{array} \right\} \leftrightarrow \begin{pmatrix} + & + & + & + \\ + & - & + & - \\ + & + & - & - \\ + & - & - & + \end{pmatrix} =: H_4
 \end{array}$$

(where we omit 1 in these elements of the Hadamard matrix)

Figure 2



$w_2^{(1)}$	++	++++	++++	++++	++++
$w_2^{(2)}$	++	+--+	+--+	+--+	+--+
$w_2^{(3)}$	++	++--	++--	++--	++--
$w_2^{(4)}$.	+--+	+--+	+--+	+--+
$w_2^{(5)}$.	++++	----	++++	----
$w_2^{(6)}$.	+--+	-+-+	+--+	-+-+
$w_2^{(7)}$		++--	--++	++--	--++
$w_2^{(8)}$		+--+	-+-+	+--+	-+-+
$w_2^{(9)}$		++++	++++	----	----
$w_2^{(10)}$		+--+	+--+	-+-+	-+-+
$w_2^{(11)}$		++--	++--	--++	--++
$w_2^{(12)}$		+--+	+--+	-+-+	-+-+
$w_2^{(13)}$		++++	----	----	++++
$w_2^{(14)}$.	+--+	-+-+	-+-+	+--+
$w_2^{(15)}$.	++--	--++	--++	++--
$w_2^{(16)}$.	+--+	-+-+	-+-+	+--+

$= H_4 \otimes H_4$

Figure 3

Now we notice that

$$P_n F := \sum_{i=1}^n \alpha_i W_i$$

is the best L_2 -approximation to a given function F from

$$\hat{M}_n := \text{span}(W_i)_{i=1}^n$$

where

$$\alpha_i := \int_{\Delta} F(p) W_i(p) dp .$$

Hence it is convergent to F if F is in L_2 , since \hat{M}_n is dense in L_2 . I.e.

Theorem 4. If $F \in L_2(\Delta)$, then $\lim_{n \rightarrow \infty} \|F - P_n F\|_2 = 0$.

Since $\hat{M}_{4^n} = M_n$ and from Theorem 3 we get the following theorem.

Theorem 5. Let $F \in C(\Delta)$, P_{4^n} be L_2 -projector onto M_{4^n} on Δ , then

$$\lim_{n \rightarrow \infty} \|F - P_{4^n} F\|_{\infty} = 0 .$$

Acknowledgement

The authors wish to express their sincere appreciation to Professor de Boor for his help.

REFERENCES

- [1] K. G. Beauchamp, Walsh Function and their Applications, Academic Press, New York, 1975.
- [2] R. B. Crittenden and L. G. Swanson, "Complete orthonormal systems generated by matrices", Proc. 9th S-E Conf. Combinatorics, Graph Theory, and Computing, 1978, 229-233.
- [3] A. Haar, "Zur Theorie der orthogonalen Funktionensysteme", Math. Ann. 69 (1910), 331-371.
- [4] M. Hall, Jr., Combinatorial Theory, Blaisdell, Waltham, Massachusetts, 1967.
- [5] H. F. Harmuth, Transmission of Information by Orthogonal Functions, second edition, Springer-Verlag, New York, Heidelberg, Berlin, 1972.
- [6] J. L. Walsh, "A closed set of normal orthogonal functions", Amer. J. Math. 45 (1923), 5-24.

YYP:DXQ:scr

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER ARC-T2K-2235	2. GOVT ACCESSION NO. AD-A103 870	3. RECIPIENT'S CATALOG NUMBER 7
4. TITLE (and Subtitle) ON THE HAAR AND WALSH SYSTEMS ON A TRIANGLE		5. TYPE OF REPORT & PERIOD COVERED Summary report, no specific reporting period
7. AUTHOR(s) Y. Y. Feng and D. X. Qi		6. PERFORMING ORG. REPORT NUMBER
8. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Madison, Wisconsin 53706		9. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041
10. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		11. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit #3 - Numerical Analysis & Computer Science
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. REPORT DATE Jul 1981
		14. NUMBER OF PAGES 18
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Haar function Uniform convergence Walsh function Triangular domain Orthogonal system Series expansion		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) In this paper we establish the Haar and Walsh systems on a triangle. These systems are complete in $L_2(\Delta)$. The uniform convergence of the Haar-Fourier series and the uniform convergence by group of the Walsh-Fourier series for any continuous function are proved.		